

Entanglement induced by nonadiabatic chaos

Hiroshi Fujisaki*

Department of chemistry, Boston University, 590 Commonwealth Ave., Boston, 02215, Massachusetts, USA

(Dated: February 1, 2008)

We investigate entanglement between electronic and nuclear degrees of freedom for a model nonadiabatic system. We find that entanglement (measured by the von Neumann entropy of the subsystem for the eigenstates) is large in a statistical sense when the system shows “nonadiabatic chaos” behavior which was found in our previous work [Phys. Rev. E **63**, 066221 (2001)]. We also discuss non-statistical behavior of the eigenstates for the regular cases.

PACS numbers: 33.80.Be, 05.45.Mt, 03.65.Ud, 03.67.-a

Quantum information processing (QIP) is one of hot topics in many branches of science [1]. One important point is how to implement a quantum computer in real systems, and many possibilities have been theoretically suggested and experimentally tested. One candidate can be molecular systems because highly excited molecules have dense quantum states, which can be manipulated by laser fields. Some quantum logic gates in such a system can be built by using optimal control theory [2], and are actually realized in a molecular system [3].

In highly excited molecules or laser-driven molecular systems, nonadiabatic transition (NT) is a rule rather than an exception [4, 5], i.e., we have to consider not only electronic or nuclear degrees of freedom (DoF), but both at the same time. In such a case, a fundamental issue related to QIP is how much (quantum) entanglement is produced between electronic and nuclear DoF in molecular systems, because entanglement is a key ingredient in QIP.

Here we investigate a two-mode-two-state (TMTS) system which has two electronic DoF and two nuclear (vibrational) DoF with a nonadiabatic coupling [6, 7]. This is a (minimum) model NT system which shows “quantum chaos” behavior, i.e., statistical properties of energy levels and eigenstates are similar to those of a random matrix system [8]. If many electronic DoF are involved, the similar system has a naive classical limit, and its dynamical property of entanglement has been already addressed in [9]. In such a case, a quantum chaological view is effective, and we can say much about a quantum system by studying its classical limit [9]. However, the situation is different and more difficult here, because the TMTS system does not have a naive classical limit due to discreteness of the electronic DoF [10], and deserves further attentions in view of entanglement.

The TMTS system in the diabatic representation is described by the following Hamiltonian:

$$\mathcal{H}_{\text{TMTS}} = \begin{pmatrix} T_{\text{kin}} + V_A & J \\ J & T_{\text{kin}} + V_B \end{pmatrix}, \quad (1)$$

where T_{kin} is the kinetic energy, V_i ($i = A, B$) is the

potential energy for state i defined by

$$T_{\text{kin}} = \frac{1}{2}(p_x^2 + p_y^2), \quad (2)$$

$$V_i = \frac{1}{2}(\omega_x^2 \xi_i^2 + \omega_y^2 \eta_i^2) + \epsilon_i \quad (i = A, B) \quad (3)$$

with

$$\xi_A = x \cos \theta - (y - a) \sin \theta, \quad (4)$$

$$\eta_A = x \sin \theta + (y - a) \cos \theta, \quad (5)$$

$$\xi_B = x \cos \theta + (y + a) \sin \theta, \quad (6)$$

$$\eta_B = -x \sin \theta + (y + a) \cos \theta. \quad (7)$$

The geometrical meaning of the parameters is shown in Fig. 1. Note that we have just used a harmonic potential for each state.




FIG. 1: A schematic representation of the TMTS system. The distance between the minima of the potential is $2a$, and the angle between the relevant crossing seam (dotted line) and the primary axis of each potential (dashed line) is θ . Inset: The perspective of the TMTS system. The potential minima are different with $\Delta\epsilon = \epsilon_B - \epsilon_A = 0.173$.

Here the Duschinsky angle θ [11] and the nonadiabatic coupling constant J are two important parameters for the system; The latter induces entanglement between electronic and vibrational DoF. We solve this Hamiltonian according to the procedure in [6], and obtain the eigen-energies and eigen-vectors. The k -th eigenvector can be written as

$$|\Phi^{(k)}\rangle = \sum_{i=1}^2 \sum_j C_{ij}^{(k)} |i\rangle |j\rangle = \sum_{i=1}^2 |\phi_i^{(k)}\rangle |i\rangle \quad (8)$$

*Electronic address: fujisaki@bu.edu

where $|1\rangle = (|A\rangle + |B\rangle)/\sqrt{2}$, $|2\rangle = (-|A\rangle + |B\rangle)/\sqrt{2}$, $|A\rangle$, $|B\rangle$ are the electronic bases for diabatic surfaces A and B , respectively, $|j\rangle$ represents 2D harmonic eigenfunctions, and $|\phi_i^{(k)}\rangle \equiv \langle i|\Phi^{(k)}\rangle = \sum_j C_{ij}^{(k)} |j\rangle$.

From this eigenvector, we can construct a reduced density operator for the electronic DoF as

$$\begin{aligned} \rho^{(k)} &= \text{Tr}_{\text{vib}}\{|\Phi^{(k)}\rangle\langle\Phi^{(k)}|\} = \begin{pmatrix} \rho_{11}^{(k)} & \rho_{12}^{(k)} \\ \rho_{21}^{(k)} & \rho_{22}^{(k)} \end{pmatrix} \\ &= \begin{pmatrix} \sum_j C_{1,j}^{(k)} (C_{1,j}^{(k)})^* & \sum_j C_{1,j}^{(k)} (C_{2,j}^{(k)})^* \\ \sum_j C_{2,j}^{(k)} (C_{1,j}^{(k)})^* & \sum_j C_{2,j}^{(k)} (C_{2,j}^{(k)})^* \end{pmatrix} \end{aligned} \quad (9)$$

where $C_{i,j}^{(k)}$ are actually all real numbers.

The measure of entanglement we choose here is the von Neumann entropy of the subsystem defined by

$$\begin{aligned} S_{\text{vN}}^{(k)} &= -\text{Tr}\{\rho^{(k)} \log \rho^{(k)}\} \\ &= -\lambda_1^{(k)} \log \lambda_1^{(k)} - \lambda_2^{(k)} \log \lambda_2^{(k)} \end{aligned} \quad (10)$$

where $\lambda_i^{(k)}$ ($i = 1, 2$) is an eigenvalue for the 2×2 matrix, Eq. (9). A note in order: the value of the entropy is the same if we use the reduced density operator for the vibrational DoF. We took the electronic DoF because the 2×2 matrix is very easy to diagonalize, and to interpret the result as shown below.

First we show the J dependence of the results fixing $\theta = \pi/6$: As we can see in Fig. 2, the entropies for the case of $J = 1.5$ assemble around its maximum $S_{\text{vN}} \simeq \log 2$, whereas those of the other cases ($J = 0.3, 7.5$) are rather broadly distributed. This condition of entanglement is very similar to that of quantum chaos behavior found in [6]: When both J and θ have “intermediate” values ($J \simeq 1$ and $\theta \simeq \pi/4$), the system shows quantum chaos behavior, i.e., the nearest neighbor spacing distribution becomes the Wigner type, Δ_3 statistics a log curve, and the amplitude distribution of the eigenstates Gaussian. To further confirm this, we show the θ dependence of the results fixing $J = 1.5$ in Fig. 3. This result also nicely corresponds to the previous condition for the quantum chaos behavior. From these results, we can conclude that, *in a statistical sense*, the condition for quantum chaos behavior to arise in the TMTS system is very similar to that for entanglement production in eigenstates to arise in the same system. This conclusion result supports the previous study [9] which uses entanglement production as an indication of quantum chaos behavior. (It is also noted that the calculation of entanglement production is rather easier than those like the nearest neighbor spacing distribution or Δ_3 statistics because there is no fitting procedure.)

However, as noticed in Fig. 2, the amount of entanglement strongly varies depending on *each* eigenstate for the cases of $J = 0.3$ and 7.5 . To consider this problem, we rewrite the entropy (entanglement) using the vibrational bases for each electronic state: $|\phi_A^{(k)}\rangle$ and $|\phi_B^{(k)}\rangle$. These states are connected to the above states $|\phi_1^{(k)}\rangle$ and $|\phi_2^{(k)}\rangle$




FIG. 2: J dependence of entanglement production measured by the von Neumann entropy as a function of the energy level number. Top: $J = 7.5$. Middle: $J = 1.5$. Bottom: $J = 0.3$. The Duschinsky angle is fixed as $\theta = \pi/6$.




FIG. 3: θ dependence of entanglement production measured by the von Neumann entropy as a function of the energy level number. Top: $\theta = \pi/3$. Middle: $\theta = \pi/6$. Bottom: $\theta = 0.0$. The nonadiabatic coupling is fixed as $J = 1.5$.

by

$$|\phi_A^{(k)}\rangle = \frac{1}{\sqrt{2}}(|\phi_1^{(k)}\rangle - |\phi_2^{(k)}\rangle), \quad (11)$$

$$|\phi_B^{(k)}\rangle = \frac{1}{\sqrt{2}}(|\phi_1^{(k)}\rangle + |\phi_2^{(k)}\rangle). \quad (12)$$




FIG. 4: $S_{\text{vN}}^{(k)}$ (+), $|\Delta P_{AB}^{(k)}|$ (*), and $|S_{AB}^{(k)}|$ (x) as a function of energy level number. (a) $J = 0.3$ (weakly nonadiabatic case). (b) $J = 7.5$ (strongly nonadiabatic case). (Lines are just for guiding eyes.) The Duschinsky angle is fixed as $\theta = \pi/6$.

Hence the density operator, Eq. (9), is represented as

$$\rho_{11}^{(k)} = \frac{1}{2} + \langle \phi_A^{(k)} | \phi_B^{(k)} \rangle \equiv \frac{1}{2} + S_{AB}^{(k)}, \quad (13)$$

$$\rho_{22}^{(k)} = \frac{1}{2} - \langle \phi_A^{(k)} | \phi_B^{(k)} \rangle \equiv \frac{1}{2} - S_{AB}^{(k)}, \quad (14)$$

$$\rho_{12}^{(k)} = \frac{1}{2} (\langle \phi_A^{(k)} | \phi_A^{(k)} \rangle - \langle \phi_B^{(k)} | \phi_B^{(k)} \rangle) \equiv \Delta P_{AB}^{(k)} \quad (15)$$

where we have introduced two new parameters: $S_{AB}^{(k)}$ is the overlap between the k -th eigenstates on surface A and B , and $\Delta P_{AB}^{(k)}$ is the half of the population difference between the k -th eigenstates on surface A and B . Using these parameters, the eigenvalues for the entropy is written as

$$\lambda_{1,2}^{(k)} = \frac{1}{2} \pm \sqrt{|S_{AB}^{(k)}|^2 + |\Delta P_{AB}^{(k)}|^2}. \quad (16)$$

From this relation, for the entropy to be large, *both* $|S_{AB}^{(k)}|$ and $|\Delta P_{AB}^{(k)}|$ should be small. We can numerically confirm this for the strongly “chaotic” case ($J = 1.5$), and this property might be derived from the random matrix theory. For less “chaotic” cases ($J = 0.3, 7.5$), the situation is different: As shown in Fig. 4, there is a strong correlation between $S_{\text{vN}}^{(k)}$ and $|\Delta P_{AB}^{(k)}|$ for the weakly nonadiabatic case ($J = 0.3$), whereas between $S_{\text{vN}}^{(k)}$ and $|S_{AB}^{(k)}|$ for the strongly nonadiabatic case ($J = 7.5$). On the other hand, $|S_{AB}^{(k)}| \simeq 0$ for the former and $|\Delta P_{AB}^{(k)}| \simeq 0$ for the latter. This is interpreted as follows: For the former, the eigenstates “reside” on diabatic surfaces A and B which are tilted each other. Thus the overlapping between the eigenstates $|S_{AB}^{(k)}|$ becomes small because the nodal patterns for the eigenstates are also tilted (see Fig. 5 in [6]). For the latter, the eigenstates “reside” on adiabatic surfaces, and the amplitudes of them on diabatic surfaces A and B are similar (see Fig. 7 in [6]), hence $|\Delta P_{AB}^{(k)}| \simeq 0$.

Let us focus on the weakly nonadiabatic case ($J = 0.3$). In the range of $k = 800$ to 820 , the lowest entangled state is 805-th, and the highest is 815-th [Fig. 4 (a)]. In Figs. 5 and 6, we show the two eigenstates on diabatic surfaces A




FIG. 5: A less entangled regular state: 805-th eigenstates on diabatic surfaces A (A) and B (B) for the weakly nonadiabatic case: $J = 0.3$. $P_i(x, y) = |\langle x, y | \phi_i \rangle|^2$ ($i = A, B$). Note that the scale for (B) is smaller than that for (A). The Duschinsky angle is fixed as $\theta = \pi/6$.




FIG. 6: A strongly entangled regular state: 815-th eigenstates on diabatic surfaces A (A) and B (B) for the weakly nonadiabatic case: $J = 0.3$. $P_i(x, y) = |\langle x, y | \phi_i \rangle|^2$ ($i = A, B$). The Duschinsky angle is fixed as $\theta = \pi/6$.

and B . As anticipated from the above argument, there is a large population difference on surface A and B for the less entangled state, whereas there is not for the strongly entangled state. The latter situation means that even a regular state can strongly entangle. Note that, albeit we desymmetrized the system with a finite $\Delta\epsilon = \epsilon_A - \epsilon_B$, we have this entangled state for the regular case. (If we

do not desymmetrize the system, i.e., $\Delta\epsilon = 0$, we easily have entangled states for both regular and chaotic cases because of the symmetry.) Thus we must be cautious to use the entanglement production as a manifestation of quantum chaotic behavior [9], though it is fine to use it in a statistical sense.

In this paper, we investigated quantum entanglement production between electronic and nuclear (vibrational) degrees of freedom for a nonadiabatic system. We found

that the condition of the entanglement in the eigenstates to appear is statistically very similar to that of quantum chaos behavior to show up. We also discussed the non-statistical behavior of the entanglement production and interpreted it using the eigenstate properties on diabatic surfaces. It will be interesting to analyze other nonadiabatic systems like Jahn-Teller molecules [12] in light of entanglement production.

-
- [1] M.A. Nielsen and I.L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
 - [2] J.P. Palao and R. Kosloff, Phys. Rev. Lett. **89**, 188301 (2002); Phys. Rev. A **68**, 062308 (2003).
 - [3] J. Vala, Z. Amitay, B. Zhang, S.R. Leone, and R. Kosloff, Phys. Rev. A **66**, 062316 (2002).
 - [4] H. Nakamura, *Nonadiabatic Transition: Concepts, Basic Theories and Applications* (World Scientific, Singapore, 2002); C. Zhu, Y. Teranishi, and H. Nakamura, Adv. Chem. Phys. **117**, 127 (2001).
 - [5] K. Takatsuka, Y. Arasaki, K. Wang, and V. McKoy, Faraday Discuss. **115**, 1 (2000); Y. Arasaki, K. Takatsuka, K. Wang, and V. McKoy, Phys. Rev. Lett. **90**, 248303 (2003).
 - [6] H. Fujisaki and K. Takatsuka, Phys. Rev. E **63**, 066221 (2001). See also H. Fujisaki and K. Takatsuka, J. Chem. Phys. **114**, 3497 (2001); H. Higuchi and K. Takatsuka, Phys. Rev. E **66**, 035203(R) (2002).
 - [7] E.J. Heller, J. Chem. Phys. **92**, 1718 (1990).
 - [8] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics* (Springer-Verlag, New York, 1990).
 - [9] K. Furuya, M.C. Nemes, and G.Q. Pellegrino, Phys. Rev. Lett. **80**, 5524 (1998).
 - [10] Note, on the other hand, that there is a sound correspondence between the TMTS system and its mapping (quasi-classical) system. See H. Fujisaki, quant-ph/0401136; Phys. Rev. E (in press).
 - [11] See J. Tang, M.T. Lee, and S.H. Lin, J. Chem. Phys. **119**, 7188 (2003), and references therein.
 - [12] H. Köppel, W. Domcke, and L.S. Cederbaum, Adv. Chem. Phys. **57**, 59 (1984); D.M. Leitner, H. Köppel, and L.S. Cederbaum, J. Chem. Phys. **104**, 434 (1996); H. Yamasaki, Y. Natsume, A. Terai, and K. Nakamura, Phys. Rev. E **68**, 046201 (2003); A.P. Hines, C.M. Dawson, R.H. McKenzie, and G.J. Milburn, quant-ph/0402016.